

Admissible partitions and the square of the Vandermonde determinant


Brian G Wybourne

Instytut Fizyki,

Uniwersytet Mikołaja Kopernika - Poland

July 2002

Dedicated to the memory of Claude Itzykson
(1938-1995)


COMMISSARIAT A L'ENERGIE ATOMIQUE - DIRECTION DES SCIENCES DE LA MATIERE
SERVICE DE PHYSIQUE THEORIQUE
CENTRE D'ETUDES DE SACLAY - ORME DES MERISIERES

Adresse Postale :
SERVICE DE PHYSIQUE THEORIQUE DE SACLAY
91191 GIF-SUR-YVETTE CEDEX, France

Téléphone : (1) 69 00 7209
Secrétariat : (1) 69 00 73 85
Fac-similé : (1) 69 00 81 20
Télex : ENRG 604641 F
Courrier électronique :

C. ITZYKSON

Mardi 1 1994

Dear Pr. Wybourne

We have just received today your very interesting contribution to the expansion of (powers of) the discriminant.

Indeed you seem to have discovered quite a striking phenomenon which deserves an explanation - Our preprint was thought - too mathematical for Nucl. Phys. B - so it is resubmitted to Journ. Mod. Phys. A published in Singapore. I hope it will be found suitable there -

The subject seems to me to be still widely open = For instance, is there a rule for the signs of the coefficients? What is the meaning of the vanishing terms you have

found? What is their general feature? Is it significant that you found them starting at $N=8$? etc...

Best wishes

Claude Itzykson

Claude Itzykson's Letter

- Indeed you seem to have discovered a striking phenomenon which deserves an explanation ...
- The subject seems to me to be still widely open = For instance, is there a rule for the signs of the coefficients?
- What is the meaning of the vanishing terms you have found?
- What is their general feature?
- Is it significant that you found them starting at $N = 8$? etc..
- Claude Itzykson March 1 1994

Expansion of the Laughlin wavefunction

- Laughlin (1983) described the fractional quantum Hall effect in terms of a wavefunction

$$\Psi_{Laughlin}^m(z_1, \dots, z_N) = \prod_{i < j}^N (z_i - z_j)^{2m+1} \exp\left(-\frac{1}{2} \sum_{i=1}^N |z_i|^2\right) \quad (1)$$

The Vandermonde alternating function in N variables is defined as

$$V(z_1, \dots, z_N) = \prod_{i < j}^N (z_i - z_j) \quad (2)$$

$$\frac{\Psi_{Laughlin}}{V} = V^{2m} = \sum_{\lambda \vdash n} c^\lambda s_\lambda \quad (3)$$

where $n = mN(N - 1)$ and the s_λ are Schur functions.

Coefficients of the expansion

- The coefficients c_λ are signed integers. Henceforth we consider the case where $m = 1$. The partitions, (λ) , indexing the Schur functions are of weight $N(N - 1)$. Algorithms exist for computing the expansions and complete results have been obtained for $N \leq 10$. For a given N the partitions are bounded by a highest partition $(2N - 2, 2N - 4, \dots, 0)$ and a lowest partition $((N - 1)^{N-1})$ with the partitions being of length N and $N - 1$. Let

$$n_k = \sum_{i=0}^k \lambda_{N-i} - k(k+1)k = 0, 1, \dots, N - 1 \quad (4)$$

Admissible Partitions

- Di Francesco *et al* define *admissible partitions* as satisfying Eq(4) with *all* $n_k \geq 0$. They computed the number of admissible partitions A_N for $N \leq 29$ and conjectured that A_N was the number of distinct partitions arising in the expansion, Eq(3), *provided none of the coefficients vanished*.
- The conjecture fails for $N \geq 8$. We find the number of admissible partitions associated with vanishing coefficients as

$$(N = 8) \quad 8, \quad (N = 9) \quad 66, \quad (N = 10) \quad 389$$

Admissible partitions whose coefficients vanish

- **Reversed partition symmetry**
- The coefficients of s_λ and s_{λ_r} are equal if

$$(\lambda_r) = (2(N - 1) - \lambda_N, \dots, 2(N - 1) - \lambda_1) \quad (5)$$

- We list the 8 partitions for $N = 8$ as reverse pairs

$$\{13 \ 11 \ 985^2 41\} \quad \{13 \ 10 \ 9^2 6531\} \quad (Q1)$$

$$\{13 \ 11 \ 9854^2 2\} \quad \{13 \ 10 \ 987531\} \quad (Q2)$$

$$\{13 \ 11 \ 976541\} \quad \{12 \ 10^2 96531\} \quad (Q3)$$

$$\{12 \ 11 \ 97^2 4^2 1\} \quad \{12 \ 10^2 7^2 532\} \quad (Q4)$$

The q -discriminant

- Let $q\mathbf{x} = (qx_1, qx_2, \dots, qx_N)$ and the q -discriminant of \mathbf{x} be

$$D_N(q; \mathbf{x}) = \prod_{1 \leq i \neq j \leq N} (x_i - qx_j) \quad (6)$$

and

$$R_N(q; \mathbf{x}) = \prod_{1 \leq i \neq j \leq N} (x_i - qx_j)(qx_i - x_j) = \sum_{\lambda} c^{\lambda}(q) s_{\lambda}(\mathbf{x}) \quad (7)$$

So that

$$V_N^2(\mathbf{x}) = \prod_{1 \leq i < j \leq N} (x_i - x_j)^2 = R_N(1; \mathbf{x}) \quad (8)$$

Introduce q -polynomials such that

$$R_N(q; \mathbf{x}) = \sum_{\lambda} c^{\lambda}(q) s_{\lambda}(\mathbf{x}) \quad (9)$$

The q -polynomials

$$R_N(q; \mathbf{x}) = \frac{(-1)^{N(N-1)/2}}{(1-q)^N} \sum_{\nu \subseteq (N-1)^N} ((-q)^{|\nu|}) + (-q)^{N^2 - |\nu|} \\ \times s_{(N-1)^N / \nu}(\mathbf{x}) s_{\nu'}(\mathbf{x})$$

Such expansions have been evaluated as polynomials in q for all admissible partitions for $N = 2 \dots 6$ with many examples for $N = 7, 8, 9$.

Some q -polynomials

N=2	[1]	q	{2}
	[-3]	$-(q^2 + q + 1)$	{1 ² }
N=3	[1]	q^3	{42}
	[-3]	$-q^2(q^2 + q + 1)$	{41 ² } + {3 ² }
	[6]	$+q(q^2 + q + 1)(q^2 + 1)$	{321}
	[-15]	$-(q^2 + q + 1)(q^4 + q^2 + q + 1)$	{2 ³ }
N=4	[1]	q^6	{642}
	[-3]	$-q^5(q^2 + q + 1)$	{641 ² } + {63 ² } + {5 ² 2}
	[6]	$+q^4(q^2 + q + 1)(q^2 + 1)$	{6321} + {543}

$$N = 8 \text{ } q\text{-polynomials where } c_\lambda(1) = 0$$

The q -polynomials for the four pairs of partitions designated earlier as $Q(1)\dots Q(4)$ are

$$\begin{aligned} Q(1) & -q^{17}(q^2 - q + 1)^2(q^2 + 1)^2(q^2 + q + 1)^5(1 - q)^4 \\ Q(2) & +q^{16}(q^2 - q + 1)^2(q^2 + 1)(q^2 + q + 1)^6(1 - q)^4 \\ Q(3) & +q^{16}(q^2 - q + 1)^2(q^2 + 1)^3(q^2 + q + 1)^5(1 - q)^4 \\ Q(4) & +q^{14}(q^2 - q + 1)^2(q^2 + q + 1)^5(1 - q)^4 \\ & \times (q^{10} + q^9 + 3q^8 + 4q^6 + q^5 + 4q^4 + 3q^2 + q + 1) \end{aligned}$$

Note the factor $(q - 1)^4$.

Some Conjectures

- If a q -polynomial is of the form $(-1)^\phi q^p Q(q)$ then under $N \rightarrow N + 1$

$$\phi \rightarrow \phi, p \rightarrow p + N, Q(q) \rightarrow Q(q), \{\lambda\} \rightarrow \{2N - 2, \lambda\}$$

- Define

$$QS(N) = \sum_{\lambda} c_{\lambda}(q)$$

then

$$QS(N) = \prod_{x=0}^{[N/2]} (-3x + 1) \prod_{x=0}^{[(N-1)/2]} (6x + 1)$$

Sum of Squares Problem

- Di Francesco et al establish the remarkable result that the sum of the squares of the coefficients of the second power of the Vandermonde with $q = 1$ is

$$\frac{(3N)!}{N!(3!)^N}$$

What is the corresponding result for the q -polynomials?

- For $N = 4$ one finds

$$\begin{aligned} & q^{24} + 6q^{23} + 22q^{22} + 58q^{21} + 128q^{20} + 242q^{19} \\ & + 418q^{18} + 646q^{17} + 929q^{16} + 1210q^{15} + 1490q^{14} \\ & + 1670q^{13} + 1760q^{12} + 1670q^{11} + 1490q^{10} + 1210q^9 \\ & + 646q^8 + 418q^6 + 242q^5 + 128q^4 + 58q^3 + 22q^2 + 6q + 1 \end{aligned}$$

Note the polynomial is symmetrical and unimodal!

References

- G V Dunne, Slater Decomposition of Laughlin States, *Int. J. Mod. Phys. B* **7**,4783 (1993)
- P Di Francesco, M Gaudin, C Itzykson and F Lesage, Laughlin's wave functions, Coulomb gases and expansions of the discriminant
Int. J. Mod. Phys. A **9**,4257 (1994)
- T Scharf, J-Y Thibon and B G Wybourne, Powers of the Vandermonde determinant and the quantum Hall effect
J Phys A:Math. Gen. **27**, 4211 (1994)
- This work has benefited from interaction with R C King and J-Y Thibon
- Support from the Polish KBN is acknowledged

- **Admissible partitions and the square of the Vandermonde determinant**
- **Brian G Wybourne**, Instytut Fizyki, Uniwersytet Mikołaja Kopernika Poland
- The Vandermonde alternating function in N variables is defined as

$$V(z_1, \dots, z_N) = \prod_{i < j}^{N} (z_i - z_j)$$

$$\frac{\Psi_{Laughlin}}{V} = V^{2m} = \sum_{\lambda \vdash n} c^\lambda s_\lambda$$

where $n = mN(N - 1)$ and the s_λ are Schur functions.

Principal Topics Considered

- Determination of the signed integers c_λ $N = 2, \dots, 10$
- Admissible partitions and zero coefficients
- The q -discriminant and q -polynomials $c_\lambda(q)$
- A conjecture

$$QS(N) = \sum_{\lambda} c_{\lambda}(q) = \prod_{x=0}^{[N/2]} (-3x + 1) \prod_{x=0}^{[(N-1)/2]} (6x + 1)$$

- Sum of squares - the q -polynomial is symmetrical and unimodal